Analysis of apoptosis in the preimplantation bovine embryo using TUNEL

A. T. Byrne¹, J. Southgate², D. R. Brison³ and H. J. Leese¹

¹Department of Biology, University of York, PO Box 373, York YO10 5YW, UK; ²ICRF Cancer Medicine Research Unit, St James's University Hospital, Leeds LS9 7TF, UK; and ³Department of Reproductive Medicine, St Mary's Hospital, Whitworth Park, Manchester M13 0JH, UK

The occurrence of cell death by apoptosis was examined in blastocyst and pre-blastocyst stage bovine embryos. Zygotes were obtained by *in vitro* maturation and *in vitro* fertilization of oocytes from abattoir derived ovaries. Two-cell to hatched blastocyst stage embryos were stained with propidium iodide to label all nuclei and by terminal deoxynucleotidyl transferase (TdT)-mediated d-UTP nick end-labelling (TUNEL) to label apoptotic nuclei, and were analysed by epifluorescent and confocal microscopy. Apoptosis was first observed at the 9–16-cell stage of development, decreasing at the morula stage before increasing at the blastocyst stage. Apoptotic dead cell index in day 7 blastocysts was negatively correlated with the total number of cells; the percentage of dead cells ranged from approximately 1 to 10% and occurred predominantly within the inner cell mass. In addition, apoptotic dead cell index was significantly higher ($P < 0.05$) in blastocysts cultured (from the two-cell stage) in the presence of 10% fetal bovine serum compared with those developed in serum-free medium. Embryos selected for early cleavage at < 29 h after fertilization and cultured together until the blastocyst stage showed a significantly lower rate of apoptosis ($P < 0.01$) compared with slower cleaving embryos.

Introduction

The presence of dead cells in the preimplantation mammalian embryo was first described by Potts and Wilson (1967). Since then it has become increasingly apparent that these cells die by apoptosis (Kerr *et al.*, 1972), a form of programmed cell death. This process ensures the regulation of cell populations and cell lineages in all major mammalian systems and has many specialized functions during development.

Light microscope and ultrastructural studies of blastocysts from many species, including mice, cows, baboons, rhesus monkeys and humans, have described dead and dying cells within the embryo (El-Shershaby and Hinchliffe, 1974; Copp, 1978; Hurst *et al.*, 1978; Enders and Schlafke, 1981; Enders *et al.*, 1982; Lopata *et al.*, 1982; Mohr and Trounson, 1982; Enders *et al.*, 1990; Hardy *et al.*, 1996). Differential labelling of trophoctoderm and inner cell mass (ICM) cells with polynucleotide specific fluorochromes has been used in mouse, human, rat, rabbit and pig embryos to describe the spatial occurrence of apoptotic cell death (Handyside and Hunter, 1986; Papaioannou and Ebert, 1988; Hardy *et al.*, 1989; Pampfer *et al.*, 1990a,b; Giles and Foote, 1995; Hardy and Handyside, 1996; Lea *et al.*, 1996; Brison and Schultz, 1997). Further assessment of embryo apoptotic cell death has been carried out using the terminal deoxynucleotidyl

transferase (TdT)-mediated d-UTP nick end-labelling technique (TUNEL). This method facilitates the analysis of nuclear DNA fragmentation *in situ*. Moreover, cells undergoing the early stages of apoptosis are preferentially labelled by the TUNEL reaction (Gavrieli *et al.*, 1992). This technique has proved useful in the quantitative measurement of cell death in mouse and human embryos in which only a few cells are present and therefore small amounts of apoptotic DNA (Jurisicova *et al.*, 1996; Pampfer *et al.*, 1997; Brison and Schultz, 1997, 1998).

In most species apoptosis is largely confined to the ICM lineage; for example, in mouse day 4 and day 5 blastocysts freshly flushed from the reproductive tract, the percentage of dead cells in the ICM is of the order of 10%, whereas in the trophoctoderm is $< 3\%$ (Hardy and Handyside, 1996). Apoptosis in the ICM may regulate this cell population, since the number of ICM cells reaches a plateau in later stage blastocysts without a decrease in mitotic cell division (Handyside and Hunter, 1986). Furthermore, redundant ICM cells that retain trophoctoderm potential after blastocyst expansion may be removed by apoptosis (Handyside and Hunter, 1986; Pierce *et al.*, 1989). Thus, it is likely that blastocyst stage apoptosis acts to eliminate cells that are damaged, are in excess, are no longer required or are developmentally incompetent. Cellular 'quality control' within the ICM is critical since this lineage forms the fetus and contains the germline.

A distinction can be made between apoptosis in mor-

phologically normal *in vitro* produced (IVP) embryos and abnormal fragmenting (that is, dying) embryos. Apoptosis has not been observed before the blastocyst stage in mouse embryos that are developing normally, although the apoptotic machinery is probably present throughout preimplantation development. When extracellular signals are blocked using the protein kinase inhibitor staurosporine, cells from mouse blastocysts and four-cell stage embryos all undergo apoptosis showing characteristic physiological and biochemical features (Weil *et al.*, 1996). Apoptosis may play a role in eliminating abnormal or defective embryos as early as the first cell cycle in mice (Jurisicova *et al.*, 1998; Warner *et al.*, 1998) and the second cell cycle in humans (Jurisicova *et al.*, 1996). This first wave of apoptosis may be related to activation of the embryonic genome, the major event of which occurs at the two-cell stage in mice and the four-cell stage in humans. Sible *et al.* (1997) obtained evidence in *Xenopus* for the presence of a maternally regulated developmental checkpoint at the stage of embryonic genome activation, which can trigger apoptosis.

There are no reports of apoptosis in the bovine preimplantation embryo. This species is of particular interest since the embryonic genome is activated relatively late in development at the 9–16-cell stage (Frei *et al.*, 1989). The effect of apoptosis on allocation of cells to the trophoblast or ICM lineages may also be important since an aberrant trophoblast:ICM cell ratio may be a contributory factor in the aetiology of the enlarged offspring syndrome in domestic animals (Thompson *et al.*, 1995). One factor that is thought to predispose to fetal oversize after transfer of IVP embryos, is the presence of serum in the culture medium. Little is known of the effect on *in vitro* derived bovine embryos of environmental factors, such as embryo culture technique and the addition of serum to medium, which may play a role in regulating cell death.

The aims of the present study were: (i) to describe the timing and distribution of apoptosis in *in vitro* derived bovine embryos; (ii) to assess the effect of culturing zygotes to the blastocyst stage in the presence of serum; and (iii) to investigate whether the selection and culturing together of embryos that have cleaved < 29 h after fertilization affects the onset and incidence of apoptosis compared with slower cleaving embryos.

A preliminary account of this work has been presented (Byrne *et al.*, 1998).

Materials and Methods

Production of embryos in vitro

Bovine ovaries were collected from an abattoir and transported to the laboratory in PBS at approximately 24–26°C. The maximum time from slaughter to oocyte collection was 8 h. Cumulus–oocyte complexes (COC) were recovered by initial ovarian dissection, followed by slashing of 1–5 mm follicles using a sterile scalpel and the subsequent release of oocytes. The COC were collected into Hepes-buffered TCM-199 medium supplemented with 10 µg heparin ml⁻¹ (Sigma, Poole) and 0.4% (w/v) BSA (Sigma).

The morphology of COC was assessed and only those with a compact non-atretic cumulus oophorus and evenly graduated cytoplasm were selected for maturation. All oocytes were washed twice in Hepes-buffered TCM-199 supplemented with 0.4% BSA, washed once in maturation medium, placed in 500 µl wells of the same medium and incubated for 24 h at 39°C under humidified 5% CO₂ in air. The medium used for maturation was TCM-199 supplemented with 10% fetal bovine serum (FBS) and 10 ng epidermal growth factor (EGF) ml⁻¹ (Sigma).

Fertilized oocytes were prepared according to the method of Thompson *et al.* (1992). Spermatozoa were prepared from frozen–thawed semen samples from a single bull of proven fertility. Semen was thawed in a waterbath at 37°C, layered on a discontinuous Percoll (Pharmacia Biotech, St Albans) gradient (45:90) and centrifuged at 2100 g for 25 min. The motile sperm pellet was washed in Hepes-buffered modified Ca²⁺-free Tyrode–albumin–sodium lactate–sodium pyruvate (TALP) medium (Lu *et al.*, 1987) and resuspended in bicarbonate-buffered fertilization TALP medium with 10 µg heparin ml⁻¹, 600 µg penicillamine ml⁻¹ and 220 µg hypotaurine ml⁻¹. The sperm concentration was adjusted to give a final concentration of 1–2 × 10⁶ ml⁻¹ when added to the fertilization wells. A 100 µl aliquot of the sperm suspension was added to each 400 µl well of fertilization TALP medium containing 50–70 COC which had been washed twice in Hepes–TALP medium and once in fertilization medium after maturation. Spermatozoa and COC were incubated together for 18–22 h under the same conditions as described for oocyte maturation.

After insemination, the remaining cumulus cells were removed by vortexing for 2 min. The putative zygotes were washed three times in Hepes-buffered synthetic oviduct fluid (SOF) and placed in groups of 15–20 in 20 µl SOF supplemented with essential and non-essential amino acids (Gibco Life Technologies, Paisley), 1 mmol glutamine l⁻¹ and 0.8% fatty-acid free BSA (SoFaaBSA; Gardner *et al.*, 1994) under mineral oil in a humidified atmosphere of 5% O₂, 5% CO₂ and 90% N₂ at 39°C.

The cultures were terminated at day 7 for all experiments on blastocysts. Only embryos with a well defined blastocoele were selected for labelling to eliminate sampling bias. Blastocyst rates were consistently 25–30% of total zygotes placed in culture.

Selection of embryo stages

Embryo stages from the two-cell to the hatched blastocyst stage were selected at the following times after fertilization (Van Soom *et al.*, 1997): two-cell, 31 h; four-cell, 38 h; five- to eight-cell, 49 h; 9–16-cell, 100 h; morula, 128 h; early blastocyst, day 6; expanded blastocyst, day 7; hatched blastocyst, day 8.

Comparison of early and later cleaving embryos

Zygotes that had completed the first cleavage division at 29 h after fertilization were moved to separate drops (15–20

in 20 µl). They were allowed to develop until day 7. Total cell counts and apoptotic cell counts were made and compared with the remaining embryos that had not completed the first cleavage division at 29 h after fertilization.

Addition of fetal bovine serum to culture medium

When required, 10% warmed FBS was added to the embryo culture medium at approximately 45 h after fertilization. Three separate batches of FBS were used for the serum comparison study: batch 1, heat-inactivated FBS (JRH Biosciences, Lenexa, KS); batch 2, heat-inactivated FBS (Sigma); batch 3, FBS (Sera-Lab Ltd, Loughborough).

TUNEL

Day 7 embryos with intact zonae pellucidae were fixed and permeabilized as described for mouse embryos (Brison and Schultz, 1996) but with an increase in the duration of permeabilization in Triton-X100 to 180 min. The embryos were washed twice in PBS–polyvinyl pyrrolidone (PBS–PVP), pH 7.4, and incubated in fluorescein-conjugated dUTP and TdT (TUNEL reagents; Boehringer Mannheim, Lewes) for 1 h at 37°C in the dark. Positive controls were incubated in DNase I (1000 U ml⁻¹; Sigma), which cleaves all DNA, for 20 min at 37°C, and washed twice before TUNEL. Negative controls were incubated in fluorescein–dUTP in the absence of TdT. After TUNEL, embryos were washed three times and counterstained with propidium iodide (50 µg ml⁻¹) after RNase treatment (50 µg RNase for 60 min at room temperature) to label all nuclei.

Fluorescence microscopy

Labelled nuclei were examined with ×40 or ×65 immersion planeofluor objectives using a Zeiss Axioplan microscope fitted with epifluorescent illumination. Three standard filter sets were used: (i) for detection of fluorescein isothiocyanate (FITC) alone (emission wavelength: 525 nm); (ii) for detection of propidium iodide alone (emission wavelength: 560 nm); and (iii) a dual filter set for detection of both fluorochromes.

Blastocysts were analysed using a Biorad MRC 600 laser-scanning confocal microscope according to the method of Brison and Schultz (1998) to confirm the distribution of TUNEL positive nuclei between the ICM and trophectoderm cell lineages.

Cell counts

Total number of cells and number of apoptotic dead cells were counted and dead cell indices were calculated by dividing the number of dead cells by the total number of cells, which included dead cells. Labelling and analysis were carried out blind on individual experimental groups of embryos.

Statistical analysis

All sets of experimental data were first tested for normality using a K-S goodness of fit test. Differences in blastocyst cell number and dead cell indices were compared using either an unpaired Student's *t* test or a one-way ANOVA with least significant difference test. Differences among the percentage of embryos developing in culture were compared by chi-squared analysis. The SPSS (6) statistical package for Windows was used for all analyses. All culture experiments were performed on between three and five separate occasions.

Results

Onset of apoptosis

All stages of development from the two-cell to the hatched blastocyst stage were analysed by TUNEL for apoptotic cell death and propidium iodide for total cell counts (Fig. 1).

Consistent with other species, there was no evidence of apoptosis from the two- to the eight-cell stage. However, apoptosis occurred in bovine embryos of normal morphology at the 9–16-cell stage, in which 30% of all embryos had at least one dead cell. The dead cell index in this group was 4–6%. Although embryos with between 9 and 16 cells were grouped together as one stage, it should be noted that apoptosis was greatest in nine-cell embryos.

A similar observation was made at the morula stage, in which almost 50% of embryos showed apoptotic cell death. Dead cell index decreased (2.0–2.5%), but increased upon commencement of blastulation (4%) and stayed relatively constant until hatching. Analysis of the percentage of embryos with either 0, 1 or ≥ 2 apoptotic nuclei revealed that the majority of blastocysts possessed at least one apoptotic nucleus.

Extent and localization of apoptotic cell death in the blastocyst

The total number of cells was compared with dead cell index for day 7 blastocysts. The overall trend indicated that the number of cells was negatively correlated with extent of apoptosis (Fig. 2). Blastocysts of a small number of cells (< 100 cells) had dead cell indices ranging from 0 to almost 10%, whereas the variability was lower in embryos of > 100 cells (0–6%).

The TUNEL reagents were capable of penetrating fixed permeabilized embryos as all nuclei of embryos treated with DNase I as a positive control showed uniform staining (data not shown). However, in embryos not treated with DNase I, a fragmented nuclear morphology characteristic of apoptosis was observed (Fig. 3a,b). On some occasions, TUNEL-negative cells of fragmented morphology were observed in the propidium iodide channel. These cells were counted as apoptotic. Furthermore, as described in rat and mouse blastocysts, not all TUNEL-positive nuclei were in the process of fragmentation (Brison and Schultz, 1997; Pampfer *et al.*, 1997).

The majority of apoptotic nuclei were located in the ICM

Fig. 1. Onset of apoptosis during development of bovine embryos. (a) Dead cell indices from the two-cell to the hatched blastocyst stage. Data are expressed as mean $\pm$ SEM. (b) Percentage of embryos in (a) with (■) 0, (▨) 1 or (□) $\geq$ 2 apoptotic nuclei. The experiment was performed at least three times using a total of 45–50 embryos per stage.

region of the blastocyst as confirmed by confocal microscopy. However, evidence of nuclear fragmentation was visible in the trophectoderm and some apoptotic bodies had been extruded into the blastocoele cavity (Fig. 3).

Effect of timing of first cleavage division

Groups of blastocysts developing from zygotes that had completed the first cleavage division at 29 h after fertilization

Fig. 2. Relationship between total number of cells and apoptotic dead cell index during development of bovine embryos. A total of 145 blastocysts produced over 6 months was analysed.

were compared with slower cleaving embryos to test for a possible relationship between rate of development and the subsequent extent of apoptosis (Table 1). Faster cleaving zygotes had significantly more cells ($P < 0.01$), significantly fewer apoptotic dead cells ($P < 0.01$) and better rates of development at the blastocyst stage ($P < 0.05$). The number of cells was normalized for the two groups and dead cell indices were compared again. This ensured that the difference in dead cell index observed was independent of the number of cells, as early cleaving embryos had more cells and, by default, a lower incidence of apoptosis (see above). A significant difference in dead cell index was still observed ($P < 0.01$).

Effect of culturing embryos in the presence or absence of serum

Groups of day 7 blastocysts that had developed in the absence of FBS had a significantly greater number of cells ($P < 0.01$) than any of the groups cultured in the presence of serum (Fig. 4a). Embryos grown without serum had a smaller dead cell index (Fig. 4b). Variation among individual serum batches was also observed; group 2 embryos (JRH Biosciences) had the smallest number of cells and the greatest dead cell index. There was no significant difference among blastocyst rates (25–28%) for the four conditions.

Discussion

The results of the present study show that, in common with other mammalian species, cell proliferation in IVP bovine embryos is associated with apoptotic cell death and that the extent of apoptosis is affected by *in vitro* culture conditions. It should be emphasised that these data refer to IVP

embryos. No measurements were made of apoptosis in *in vivo* embryos and it is possible that the results do not reflect the situation *in vivo*. Further work is required to clarify this matter.

Apoptosis was first observed at the 9–16-cell stage, in embryos of normal morphology. This was an unexpected observation, since in mouse embryos apoptosis does not occur in eight-cell embryos or morulae derived either *in vivo* or *in vitro* (Handyside and Hunter, 1986; Brison and Schultz, 1997). Similarly, Jurisicova *et al.* (1996) reported that in normal human embryos, from the two-cell to the un-compacted morula stage, TUNEL analysis combined with DAPI staining (to assess chromatin condensation and nuclear shape) was negative with regard to necrosis and apoptosis. However, Jurisicova *et al.* (1996) reported that from the two-cell stage onwards, fragmenting human embryos were dying by apoptosis. (The term fragmenting is used here to describe dead or dying embryos. It does not refer to the fragmentation of DNA as occurs in apoptosis but to the morphology of the whole embryo.) This was also reported in mouse embryo development, in which a wave of fragmentation was associated with apoptosis at the one-cell stage (Jurisicova *et al.*, 1998). In this case, apoptosis may be related to activation of the embryonic genome. In mice and humans, the major events take place at the two- and four-cell stage, respectively. However, in mice, activation of the embryonic genome is initiated late in the first cell cycle (Schultz 1993), co-incident with the first observation of apoptosis in this species. The function of this wave of apoptosis may be to kill embryos that do not undergo appropriate activation of the embryonic genome. However, this does not explain the early expression of apoptosis in cows, as the embryos were neither fragmenting nor abnormal. The simultaneous activation of apoptotic cell death and the embryonic genome at the 9–16-cell stage may, when necessary, eliminate non-viable cells in the developing embryo. This proposal is supported by the high variation in the extent of apoptosis at this time. It would be of interest to correlate apoptosis with transcription of early markers of activation of the embryonic genome in individual blastomeres.

A marked increase in the incidence of apoptosis was apparent after the onset of blastulation. Most embryos (90–100%) showed some cell death from the early to the hatched blastocyst stage compared with earlier stages. Apoptotic index was inversely related to the number of blastocyst cells, indicating that after initial activation, cell death decreased with blastocyst expansion. Most apoptosis appeared to be present in the ICM. These observations are similar to those in mouse and human blastocysts (Hardy, 1997) and are consistent with apoptosis serving a physiological histogenetic function (Handyside and Hunter, 1986). As murine blastocyst development progresses, ICM cells lose their ability to regenerate trophectoderm (Handyside, 1978). It is possible that in the late stage ICM, apoptosis reaches a certain threshold to prevent the formation of ectopic trophectoderm in the incipient germ layers (Pierce *et al.*, 1989; Parchment, 1991, 1993). In the present study, in IVP bovine blastocysts, apoptosis was observed predominantly in the ICM and not in the

Fig. 3. Detection of apoptotic and all nuclei in bovine embryos by TUNEL (fluorescein isothiocyanate-conjugated dUTP; green channel) and propidium iodide (red channel), respectively. (a) Day 7 expanded blastocyst showing labelling in both channels. (b) Day 8 hatched blastocyst showing apoptotic nuclei in the inner cell mass (ICM) and also extruded into the blastocoele. (c) Confocal section through day 7 blastocyst confirming most of the apoptotic nuclei are in the ICM. (d,e) 18-cell embryo shown in the red (d) and green (e) channels showing apoptotic nuclear morphology. Scale bars represent (a,d,e) 25 μ m and (b,c) 15 μ m.

Table 1. Number of cells, dead cell index and blastocyst rates for day 7 bovine embryos that completed the first cleavage division at 29 h after fertilization (early cleaving) compared with slower cleaving embryos (late cleaving)

Parameter	Early cleaving*	Late cleaving**
Number of cells (mean $\pm$ SEM)	107.7 $\pm$ 5.3	85.65 $\pm$ 4.3 ^a
Dead cell index (%)	3.0 $\pm$ 0.3	14.9 $\pm$ 0.4 ^b
Normalized dead cell index (%)	3.5 $\pm$ 0.3	4.5 $\pm$ 0.4 ^c
Blastocyst rates (%)	24.2	11.2 ^d

*Measured on four occasions; $n = 57$, mean $\pm$ SEM.

**Measured on four occasions; $n = 40$, mean $\pm$ SEM.

^{abc} $P < 0.01$; ^d $P < 0.05$.

trophectoderm. IVP blastocysts from other species appear to lack such regulation; apoptosis is distributed randomly in both ICM and trophectoderm (humans: Hardy *et al.*, 1989; mice: Jurisicova *et al.*, 1998). Cultured blastocysts lacking the transforming growth factor α (TGF- α) gene also show extensive apoptosis outside the ICM (Brison and Schultz, 1998). The mechanism for regulation of apoptosis in the trophectoderm is currently unclear.

Apoptosis may also be related to 'embryo quality'. Although it is not possible to measure the number of blastocyst cells before embryo transfer, it is thought that embryos with a large number of cells are more likely to implant and give rise to live offspring (Van Soom *et al.*, 1997). The results of the present study show that blastocysts consisting of a small number of cells and hence possibly of reduced developmental potential, have a high and variable incidence of apoptosis. It has been suggested that when programmed cell death reaches a certain threshold, it is detrimental to further development (Jurisicova *et al.*, 1996; Hardy, 1997) thus eliminating potentially non-viable offspring. Similarly, in humans and cows, the timing of the first cleavage division is considered to be a useful non-invasive method for the selection of good quality embryos for transfer (Shoukir *et al.*, 1997; Van Soom *et al.*, 1997). For example, in humans, significantly more clinical pregnancies were obtained from embryos that had completed the first cleavage division at 25 h after fertilization (Shoukir *et al.*, 1997). Comparison of the extent of apoptosis between early and late cleaving bovine embryos also indicates that in the late cleaving and thus less viable embryos, a non-physiological incidence of cell death may result in decreased developmental competence.

Addition of serum to the culture medium has an effect on bovine and ovine embryo morphology (Gardner *et al.*, 1994), ultrastructure (Dorland *et al.*, 1994; Thompson *et al.*, 1995), metabolism (Thompson *et al.*, 1996) and post-implantation development (Walker *et al.*, 1992). It has also been implicated in the enlarged offspring syndrome (Thompson *et al.*, 1995). Culture in the presence of serum affects the developmental kinetics of embryos leading to premature blastulation (Walker *et al.*, 1992a,b). Contrary to the findings of Van Langendonck *et al.* (1997), the results of the present study show for the first time that the total number of cells is decreased and the incidence of apoptosis is increased in the presence of FBS. Furthermore, the extent to which this occurs

Fig. 4. Effect of addition of 10% fetal bovine serum (FBS) on (a) total number of cells and (b) dead cell index during development of bovine embryos. -FBS ($n = 29$), control; FBS 1 ($n = 53$), heat-inactivated FBS (JRH Biosciences, Lenexa, KS); FBS 2 ($n = 22$), heat-inactivated FBS (Sigma); FBS 3 ($n = 41$), FBS (Sera-Lab Ltd, Loughborough). The experiment was performed over 13 individual culture weeks and was repeated a minimum of three times for each condition. * $P < 0.05$, ** $P < 0.01$ compared with -FBS control.

varies with individual batches of serum, a fact that emphasises the problems of using complex media. Heat inactivation of FBS played no role in determining rates of apoptosis. Further clarification of the effect of FBS on the total number of cells is required. FBS contains a variety of constituents, such as peptide growth factors, which can act as survival factors in embryos (Brison and Schultz, 1997, 1998), but may also contain factors that can trigger apoptosis, such as the cytokine, tumour necrosis factor (TNF).

It has been suggested that embryo culture in the presence of serum affects the distribution of cells between the trophectoderm and the ICM. An increase in the

trophectoderm:ICM cell ratio may lead to an imbalance in placental and fetal size and observed pathological postnatal phenotype (Leese *et al.*, 1998). Alternatively, apoptosis may act to maintain ICM fidelity by ridding the embryo of aberrant cells, thus any alteration in the extent of cell death could potentially affect the future development of the fetus.

This work was supported by the European Union and a studentship from the University of York to A. T. Byrne. The authors wish to thank E. Ferguson and C. Hall for their help with *in vitro* embryo production.

References

- Brison DR and Schultz RM (1996) RT-PCR-based method to localise the spatial expression of genes in the mouse blastocyst *Molecular Reproduction and Development* **44** 171–178
- Brison DR and Schultz RM (1997) Apoptosis during mouse blastocyst formation: evidence for a role for survival factors including TGF- α *Biology of Reproduction* **56** 1088–1096
- Brison DR and Schultz RM (1998) Increased incidence of apoptosis in transforming growth factor α -deficient mouse blastocysts *Biology of Reproduction* **59** 136–144
- Byrne AT, Brison DR, Southgate J and Leese HJ (1998) *Journal of Reproduction and Fertility Abstract Series* **21** 18
- Copp AJ (1978) Interaction between inner cell mass and trophoctoderm of the mouse blastocyst *Journal of Embryology and Experimental Morphology* **48** 109–125
- Dorland M, Gardner DK and Trounson AO (1994) Serum in synthetic oviduct fluid causes mitochondrial degeneration in ovine embryos *Journal of Reproduction and Fertility Abstract Series* **13** 70
- El-Shershaby AM and Hinchliffe AR (1974) Cell redundancy in the zona-intact preimplantation mouse blastocyst: a light and microscope study of dead cells and their fate *Journal of Embryology and Experimental Morphology* **31** 643–654
- Enders AC and Schlafke S (1981) Differentiation of the blastocyst of the rhesus monkey *American Journal of Anatomy* **162** 1–21
- Enders AC, Hendrickx AG and Binkerd PE (1982) Abnormal development of blastocysts and blastomeres in the rhesus monkey *Biology of Reproduction* **26** 353–366
- Enders AC, Lantz KC and Schlafke S (1990) Differentiation of the inner cell mass of the baboon blastocyst *Anatomical Records* **226** 237–248
- Frei RE, Schultz GA and Church RB (1989) Qualitative and quantitative changes in protein synthesis occur at the 8–16 cell stage of embryogenesis in the cow *Journal of Reproduction and Fertility* **86** 637–641
- Gardner DK, Lane M, Spitzer A and Batt PA (1994) Enhanced rates of cleavage and development for sheep zygotes cultured to the blastocyst stage *in vitro* in the absence of serum and somatic cells: amino acids, vitamins and culturing embryos in groups stimulate development *Biology of Reproduction* **50** 390–400
- Gavrieli Y, Sherman Y and Ben-Sasson SA (1992) Identification of programmed cell death *in situ* via specific labelling of DNA fragmentation *Journal of Cell Biology* **119** 493–501
- Giles JR and Foote RH (1995) Rabbit blastocyst: allocation of cells to the inner cell mass and trophoctoderm *Molecular Reproduction and Development* **41** 204–211
- Handyside AH (1978) Time of commitment of inside cells isolated from preimplantation mouse embryos *Journal of Embryology and Experimental Morphology* **45** 37–53
- Handyside AH and Hunter S (1986) Cell division in the mouse blastocyst before implantation *Roux's Archives of Developmental Biology* **195** 519–526
- Hardy K (1997) Cell death in the mammalian blastocyst *Molecular Human Reproduction* **3** 919–925
- Hardy K and Handyside AH (1996) Metabolism and cell allocation during parthenogenetic mouse preimplantation development *Molecular Reproduction and Development* **43** 313–322
- Hardy K, Handyside AH and Winston RML (1989) The human blastocyst: cell number, death and allocation during late preimplantation development *in vitro*. *Development* **107** 597–604
- Hardy K, Warner A, Winston RML and Becker DL (1996) Expression of intercellular junctions during preimplantation development of the human embryo *Molecular Human Reproduction* **2** 621–632
- Hurst PR, Jeffries K, Eckstein P and Wheeler AG (1978) An ultrastructural study of preimplantation uterine embryos of the rhesus monkey *Journal of Anatomy* **126** 209–220
- Juriscova A, Varmuza S and Casper RF (1996) Programmed cell death and human embryo fragmentation *Molecular Human Reproduction* **2** 93–98
- Juriscova A, Rogers I, Fasciani A, Casper RF and Varmuza S (1998) Effects of maternal age and conditions of fertilization on programmed cell death during murine preimplantation embryo development *Molecular Human Reproduction* **4** 139–145
- Kerr JF, Wyllie AH and Currie AR (1972) Apoptosis: a basic biological phenomenon with wide ranging implications in tissue kinetics *British Journal of Cancer* **26** 239–257
- Lea RG, McCracken JE, McIntyre SS, Smith W and Baird JD (1996) Disturbed development of the preimplantation embryo in the insulin-dependant diabetic BB/E rat *Diabetes* **45** 1463–1470
- Leese HJ, Donnay I and Thompson JG (1998) Human assisted conception: a cautionary tale. Lessons from domestic animals *Human Reproduction* **13** Supplement 4 184–202
- Lopata A, Kohlman DJ and Kellow GN (1982) The fine structure of human blastocysts developed in culture. In *Embryonic Development Part B: Cellular Aspects* pp 69–85 Eds R Alan. Liss, New York
- Lu KH, Gordon I, Gallager M and McGovern H (1987) Pregnancy established in cattle by transfer of embryos derived from *in vitro* fertilisation of oocytes matured *in vitro*. *Veterinary Records* **121** 259–260
- Mohr LR and Trounson AO (1982) Comparative ultrastructure of hatched human, mouse and bovine blastocysts *Journal of Reproduction and Fertility* **66** 499–504
- Pampfer S, De Hertogh R, Vanderheyden I, Michiels B and Vercheval M (1990a) Decreased inner cell mass proportion in blastocysts from diabetic rats *Diabetes* **102** 793–803
- Pampfer S, Vanderheyden I, Michiels B and De Hertogh R (1990b) Cell allocation to the inner cell mass and the trophoctoderm in rat embryos during *in vivo* preimplantation development *Roux's Archives of Developmental Biology* **198** 257–263
- Pampfer S, Vanderheyden I, McCracken J, Vasela J and De Hertogh R (1997) Increased cell death in rat blastocysts exposed to maternal diabetes *in utero* and to high glucose or tumour necrosis factor- α *in vitro*. *Development* **124** 4827–4836
- Papaioannou VE and Ebert KM (1988) The preimplantation pig embryo: cell number and allocation to trophoctoderm and inner cell mass of the blastocyst *in vivo* and *in vitro*. *Development* **102** 793–803
- Parchment RE (1991) Programmed cell death (apoptosis) in murine blastocysts: extracellular free radicals, polyamines and other cytotoxic agents *In Vivo* **5** 493–500
- Parchment RE (1993) The implications of a unified theory of programmed cell death, polyamines, oxyradicals, and histogenesis in the embryo *International Journal of Developmental Biology* **37** 75–83
- Pierce GB, Lewellyn AL and Parchment RE (1989) Mechanism of programmed cell death in the blastocyst *Proceedings National Academy of Science USA* **86** 3654–3658
- Potts DM and Wilson IB (1967) The preimplantation conceptus of the mouse at 90 hours post-coitum *Journal of Anatomy* **102** 1–11
- Schultz RM (1993) Regulation of zygotic gene activation in the mouse *Bioessays* **15** 531–538
- Shoukir Y, Campana A, Farley T and Sakkas D (1997) Early cleavage of *in vitro* fertilized human embryos to the 2-cell stage: a novel indicator of embryo quality and viability *Human Reproduction* **12** 1531–1536
- Sible JC, Anderson JA, Lewellyn AL and Maller JL (1997) Zygotic transcription is required to block a maternal programme of apoptosis in *Xenopus* embryos *Developmental Biology* **189** 335–346
- Thompson JG, Pugh PA, Tervit HR and Berg DK (1992) *In vitro* production of domestic animals: advances made at Ruakura, New Zealand *Australian Biotechnology* **2** 280–287
- Thompson JG, Gardner DK, Pugh PA, McMillan WH and Tervit HR (1995) Lamb birth weight is affected by culture system utilised during *in vitro* pre-elongation development of ovine embryos *Biology of Reproduction* **53** 1385–1391
- Thompson JG, Bell CS and Tervit HR (1996) Utilisation of [1-14-C]-pyruvate by day-6 ovine blastocysts derived from *in vivo* or *in vitro* sources *Proceedings of the International Congress for Animal Reproduction, Sydney* **3** 22–24

- Van Langendonckt A, Donnay I, Schuurbiers N, Auquier P, Carolan C, Massip A and Dessy F (1997) Effects of supplementation with fetal calf serum on development of bovine embryos in synthetic oviductal fluid medium *Journal of Reproduction and Fertility* **109** 87–93
- Van Soom A, Ysebaert M and De Kruif A (1997) Relationship between timing of development, morula morphology and cell allocation to the inner cell mass and trophoctoderm in *in vitro* produced embryos *Molecular Reproduction and Development* **47** 47–56
- Walker SK, Heard TM and Seamark RF (1992a) *In vitro* culture of sheep embryos without co-culture: successes and perspectives *Theriogenology* **37** 111–126
- Walker SK, Heard TM, Bee CA, Frensham AB, Warnes DM and Seamark RF (1992b) Culture of embryos from farm animals. In *Embryonic Development and Manipulation in Animal Production* pp 111–120 Eds A Lauria and F Gandolfi. Portland Press, London
- Warner CM, McElhinny AS, Wu L, Cieluch C, Ke X, Cao W, Tang C and Exley GE (1998) Role of the Ped gene and apoptosis in control of preimplantation development *Journal of Assisted Reproduction and Genetics* **15** 331–337
- Weil M, Jacobson MD, Coles HS, Davies TJ, Gardner RL, Raff KD and Raff KC (1996) Constitutive expression of the machinery for programmed cell death *Journal of Cell Biology* **133** 1053–1059